



Platelet Rich Plasma a New Prospective in Treatment of Recalcitrant Erosive Lichen Planus – A Case Report

**Sameeulla Shaik^{1*}, P. Naga Jyothi¹, B. Vijay Kumar¹, S. V. Suman¹,
K. Sai Praveen¹ and Mercy Sravanthi¹**

¹*Department of Oral Medicine and Radiology, CKS Theja Institute of Dental Sciences and Research, Tirupathi, AP -517501, India.*

Authors' contributions

This work was carried out in collaboration among all authors. Authors BVK and PNJ designed the study. Author KSP performed the statistical analysis. Authors SS and PNJ wrote the protocol, wrote the first draft of the manuscript and managed the analyses of the study. Authors SVS and MS managed the literature searches. All authors read and approved the final manuscript.

Article Information

Editor(s):

(1) Dr. Hamit Serdar Çöttert, Professor, Department of Prosthodontics, Faculty of Dentistry, Ege University, Izmir, Turkey.

Reviewers:

(1) Sami El Toum, Lebanese University, Lebanon.

(2) Zile Rubab, Ziauddin University, Pakistan.

(3) Fedora Della Vella, University of Bari Aldo Moro, Italy.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/53790>

Case Report

Received 02 November 2019

Accepted 07 January 2020

Published 15 January 2020

ABSTRACT

Oral lichen planus(OLP) is a chronic disease that worsens the quality of life of patients with risk of malignant potential. Treatment of OLP remains a significant challenge despite the recent advances in understanding the immunopathogenesis. The properties of Platelet-rich plasma (PRP) because of its growth factors, such as epidermal growth factor (EGF), fibroblast growth factor (FGF), Transforming growth factor(TGF β1) and keratinocyte growth factor (KGF) suggested its application in clinical practice for treatment of OLP patients that do not respond to conventional therapy. Hereby we report a case of erosive OLP resistant to conventional therapy efficiently treated with PRP.

Keywords: Oral Lichen Planus; platelet rich plasma; fibroblast growth factor.

**Corresponding author: Email: drsamibds@gmail.com;*

1. INTRODUCTION

Oral lichen planus(OLP) is a relatively common chronic inflammatory disease with risk of malignant transformation. Clinically OLP can present six different patterns: papule, reticular, plaque, erythematous, erosive and bullous, each showing specific characteristics. Severe forms of OLP, i.e., erosive and ulcerative forms, worsen the quality of life of the patients with symptoms like burning sensation, pain, and risk of malignant transformation [1]. Exact etiology of lichen planus is still unclear, it hypothesized to be due to autoimmunity with deficient antigen-specific immunosuppression and lack of transforming growth factor (TGF β 1) [2]. Most of the cases of OLP tend to be more persistent and more resistant to treatment despite of many treatment modalities like topical steroids (considered as first line of treatment) [3], systemic steroids, immunosuppressants, natural agents like curcumin, aloe vera, vitamin A, biomodulation using laser [4]. This further encourages the newer treatment modalities for the treatment of OLP.

Recent literature has focused on the application of PRP, a plasma concentrate of the patient's blood that predominantly contains platelets as an adjuvant treatment modality for many medical conditions. Activated platelets in PRP release different growth factors that contribute to cell migration, proliferation, differentiation, angiogenesis, removal of tissue debris, and regeneration of the appropriate type of tissues [5]. Hereby we report a case of erosive OLP refractory to conventional therapy treated with autologous PRP.

2. CASE REPORT

A 19-year-old female patient visited a local dentist with a chief complaint of burning sensation one year back. She underwent clinical examination, incisional biopsy, and was diagnosed as OLP. She was advised application of topical steroid (Triamcnenolone acetonide 0.1%) four times a day for one-month following which the lesion regressed. Following discontinuation of medication, the lesion

reappeared. On her further, follow up visit, she was advised systemic steroid (prednisone 40 mg once daily) for two months followed with tapering dose. But her complaint remained unaltered and was referred to our institution. On examination of oral cavity, erosive lesion with white keratotic striae (wickham's striae) was found bilaterally on the right and left buccal mucosa. Gingiva and tongue was normal. Skin and genital involvement was not seen. Conventional hematologic and biochemical tests were performed, and no abnormality was detected. Incisional biopsy was taken from right buccal mucosa under local anesthesia using 2% lidocaine and was sent for histopathological examination. Based on the clinical picture and further confirmation with histopathology, we arrived at a diagnosis of erosive OLP. Considering the recurrent nature of OLP, intralesional injections of autologous PRP was administered because of its promising results with multiple growth factors.

Autologous PRP was prepared as per the standardized procedure [6]. Intra-lesional PRP injection was given after a field block with lidocaine 2% anesthetic solution. 0.5 ml of PRP was injected per 1 cm² of ulcerated mucosa using a 25 gauge needle [6]. Patient was advised to stop any other medication, and PRP injections were administered once in a week up to 4 consecutive weeks. Following termination of intralesional injections following visits were placed at an interval of 1 week, one month, 3rd months, and six months. The patient was advised to report if there were any flare-up episodes of the lesion. On follow up visits patient symptoms were analyzed using VAS score and clinical presentation was evaluated for change in size and inflammation.

The result was satisfactory with a significant reduction in patient symptoms (Table 1) and clinical presentation of the lesion after one week and complete regression of the lesion after 4th week was observed in terms of size and inflammation, and post inflammatory pigmentation was seen (Figs.1-3), On follow up visit, no recurrence of the lesion was observed even after six months follow up.

Table 1. Reduction in VAS Score

	Before treatment	After 1 st week	After 2 nd week	After 3 rd week	After 4 th week
VAS	7	5	3	1	0



Fig. 1. Appearance of lesion bilaterally during first visit



Fig. 2. Appearance of lesion after one month follow up



Fig. 3. Appearance of lesion 6 months of follow up showing post inflammatory pigmentation

3. DISCUSSION

Lichen planus is a relatively common disorder of the stratified squamous epithelia [1]. Skin lesions in lichen planus usually resolve within one or two years, whereas in OLP it may persist for 20 years or more [7]. The oral mucosa is commonly involved and maybe the only site of involvement and is estimated to affect 0.5 to 2% of population [3]. Clinically OLP can present six different patterns: papule, reticular, plaque, atrophic, erosive and bullous, each showing specific characteristics and appearing in either isolated or associated forms. Among all phenotypes erosive form being less frequent, presents greater clinical significance as the lesions are usually symptomatic, ranging from minimal discomfort to episodes of severe pain as well as malignant potential (0.2% to 1.4%) [8] and tends to be more resistant to treatment [1].

OLP is hypothesized to be an autoimmune disease with deficient antigen-specific immunosuppression in OLP with lack of TGF β 1, which may predispose to autoimmune lymphocytic inflammation, which accounts for its chronicity [2].

Topical steroids considered as first choice of agents and for severe cases systemic steroids are considered as treatment of choice, but prolonged use should be avoided because of its adverse effects like mucosal atrophy, candidiasis, adrenal suppression, gastrointestinal upset, hypertension, hyperglycemia [9]. Immunosuppressive agents affect the severity and progression of OLP but theoretically they could also trigger malignant transformation. Despite many other treatment modalities including natural agents like curcumin, aloe vera, vitamin A, biomodulation using laser still OLP tends to be resistant and recurrence is common [3,4].

PRP is a plasma concentrate of the patient blood that predominantly contains platelets. Activated platelets release different growth factors, including transforming growth factor β (TGF β 1 and TGF β 2 isomeric) that contribute to cell migration, proliferation, differentiation, angiogenesis, and removal of tissue debris and regeneration of the appropriate type of tissues [5]. It is reported that the total amount of growth factors is released around one hour after platelet activation, while 70% of the growth factors are released 10 min after the activation of thrombocytes; the autologous use is free from

immune reaction or allergy, and severe side effects are rare in the literature [10].

Lore et al. [11] in a pilot study compared the effect of PRP gel with cyclosporine mouthwash and retinoic acid lotion in different OLP phenotypes. They concluded that PRP is to be used in erosive type, which proved to be effective when applied once weekly. Mehdat EL-Komy et al. [12] in their pilot study on resistant oral erosion of pemphigus vulgaris treated with weekly injections of PRP reported improvement in healing of oral lesions and decrease in pain and discomfort associated with oral ulcers and erosions.

In the present case results was satisfactory. After 1 week follow up, reduction in size was evident and after 4 weeks follow up lesion disappeared completely. On a month, 3 months, 6months follow up no signs of recurrence were seen.

4. CONCLUSION

PRP showed to be effective in decreasing the symptoms and improvement in clinical signs of OLP, which was resistant to conventional therapy. Further investigations by larger randomized control trials and longer follow up are required to approve this method as a standard treatment modality for patients with resistant erosive OLP.

CONSENT

As per international standard, patient's consent has been collected and preserved by the authors.

ETHICAL APPROVAL

As per international standard, written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Scully C, Beyli M, Ferreiro MC, et al. Update on oral lichen planus: Etiopathogenesis and management. Crit Rev Oral Biol Med. 1998;9:86–122.
2. Roopashree MR, Gondhalekar RV, Shashikanth MC, George J, Thippeswamy

- SH, Shukla A. Pathogenesis of oral lichen planus – A review. *J Oral Pathol Med.* 2010;39:729–734.
3. Gupta S, Ghosh S, Gupta S. Interventions for the management of oral lichen planus: A review of the conventional and novel therapies. *Oral Dis.* 2017;23:1029–1042.
 4. Fornaini C. LLLT in the Symptomatic Treatment of Oral Lichen Planus. *Laser Ther.* 2012;21:51–53.
 5. Pietrzak WS, Eppley BL. Scientific Foundations Platelet Rich Plasma: Biology and New Technology. *The Journal of Craniofacial Surgery.* 2005;16(6):1043–54.
 6. Merigo E, Oppici A, Parlato A, Cella L, Clini F, Fontana M, Fornaini C. Platelet-Rich Plasma (PRP) Rinses for the Treatment of Non-Responding Oral Lichen Planus: A Case Report. *Biomedicine.* 2018;6(1):1-4.
 7. Eisenberg E. Oral lichen planus: a benign lesion. *J Oral Maxillofac Surg.* 2000; 58(11):1278-1285.
 8. Giuliani M, Troiano G, Cordaro M, Corsalini M, Gioco G, Lo Muzio L, Pignatelli P, Lajolo C. Rate of malignant transformation of oral lichen planus: A systematic review. *Oral Dis.* 2019;25(3): 693-709.
 9. Passeron T, Zakaria W, Ostovari N, Montoux F, Lacour JP, Ortonne JP. Treatment of erosive lichen planus by 308nm excimer laser. *Lasers Surg Med.* 2004;34 (4):205-209.
 10. Raeissadat SA, Babae M, Rayegani SM, Hashemi Z, Hamidieh AA, Mojangi P, Fouladi Vanda H. An overview of platelet products (PRP, PRGF, PRF, etc.) in the Iranian studies. *Future Sci.* 2017;3(4):231 34.
 11. Loré B, Saraceno R, Poladas G, Fida M, Khoury C, Arcuri C, Magnato R. Oral lichen planus: therapy and phenotype. *G Ital Dermatol Venereol.* 2018;153(4):459-463.
 12. El-Komy MHM, Hassan AS, Raheem HMA, Doss SS, El-Kaliouby M, Saleh NA, Saleh MA. Platelet-rich plasma for resistant oral erosions of pemphigus vulgaris: A pilot study. *Wound Repair and Regeneration.* 2015;23(6):953–955.

© 2020 Shaik et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/53790>